

Some notions and properties on Convergence of functions

Point wise convergence

A sequence of functions $\{f_n\}_{n \geq 0}$ defined on a set S converges pointwise to a function f defined on S if

$$\lim_{n \rightarrow \infty} f_n(x) = f(x) \text{ holds } \forall x \in S.$$

In other words,
 f_n converges pointwise to f on S if $\forall x \in S$
and $\forall \varepsilon > 0 \exists N \in \mathbb{N}$ / $\forall n \geq N$
 $|f_n(x) - f(x)| < \varepsilon$. Here N depends on
both ε and x .


Example: Consider the graph of a continuous function
 $f_n(x) = x^n$ over $(-1, 1]$

On this set

$$f(x) = \begin{cases} 0 & -1 < x < 1 \\ 1 & x = 1 \end{cases}$$

\Rightarrow The limiting function i.e., $f(x)$ is discontinuous
Point wise limit of a cont. function need not be continuous.

Example : Consider a sequence of piecewise linear functions "tent" functions



$$f_n(x) = \begin{cases} -n^2x & 0 \leq x < \frac{1}{n} \\ 2n - n^2x & \frac{1}{n} \leq x \leq \frac{2}{n} \\ 0 & \frac{2}{n} \leq x \leq 1 \end{cases}$$

Let us examine if $f_n(x)$ goes to zero pointwise on $[0, 1]$.
 If $x \in (0, 1]$, $f_n(x) = 0 \neq n$ $\forall x > \frac{2}{n}$
 If $x = 0$, $f_n(0) = 0 \neq n$
 Observe : $\int_0^1 f_n(x) dx = 1 \neq n$ ($\because \frac{1}{2} \cdot \frac{2}{n} \cdot n = 1$)
 $I = \lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx \neq \int_0^1 \lim_{n \rightarrow \infty} f_n(x) dx = \int_0^1 0 \cdot dx = 0$

UNIFORM CONVERGENCE

A sequence $\{f_n\}_{n \geq 0}$ defined on a set S converges uniformly to a function f if for every $\varepsilon > 0$

$$\exists N \in \mathbb{N} \quad / \quad n \geq N \quad |f_n(x) - f(x)| < \varepsilon$$

holds $\forall x \in S$

i.e., $\forall \varepsilon > 0 \quad \exists N \in \mathbb{N} \quad / \quad \forall n \geq N, \forall x \in S$

$$|f_n(x) - f(x)| < \varepsilon \quad \text{i.e., } \frac{N \text{ depends on } \varepsilon}{\text{but not on } x}$$

NOTE: If f_n converges to f uniformly on S , then
 f_n converges to f pointwise as well

Example: Let us examine if
 $\{f_n := \frac{nx^2+1}{nx+1}\}$ is uniformly convergent over $[1, 3]$

First, let us take the "pointwise limit"

$$\lim_{n \rightarrow \infty} \frac{nx^2+1}{nx+1} = \lim_{n \rightarrow \infty} \frac{x^2 + 1/n}{x + 1/n} = x$$

i.e., f_n converges to x pointwise over $[1, 3]$

Let us examine uniform convergence.

Consider $|f_n(x) - f(x)| = |f_n(x) - x|$

$$\left| \frac{nx^2 + 1}{nx + 1} - x \right| = \left| \frac{1 - x}{nx + 1} \right| \leq \frac{1 + |x|}{nx + 1}$$

Over $[1, 3]$, $\frac{1 + |x|}{nx + 1}$ can be upper bounded

to $\frac{4}{n+1}$ if $x \in [1, 3]$

If $\varepsilon > 0$ is chosen $\exists N / n \geq N$

$$\frac{4}{n+1} < \varepsilon$$

$$\Rightarrow n \geq N$$

$$\left| f_n(x) - f(x) \right| < \varepsilon \quad \forall x \in [1, 3]$$

This proves

UNIFORM CONVERGENCE

Applications

We know that the Fourier series for a 2π periodic function is

$$\frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(kx) + b_k \sin(kx)$$

In functional series form, the above can be written as

$$\sum_{k=0}^{\infty} s_k(x) = \lim_{N \rightarrow \infty} \sum_{k=0}^N s_k(x)$$

If it is also obvious that different values of x can give different limits.

Before we go further, let us recall the basic defns
of supremum & infimum.

Defn : Let $S \subset \mathbb{R}$. The supremum of S
denoted by $\sup S$ is the smallest number
 $a \in \mathbb{R} / x \leq a \nexists x \in S$
 $\sup S = \min \{ a \in \mathbb{R} : x \leq a \nexists x \in S \}$

III by infimum of $b \in \mathbb{R}$ such that $x \geq b \forall x \in S$
is the largest number
 $\inf S = \max \left\{ b \in \mathbb{R} : x \geq b \forall x \in S \right\}$

Let us define for a real valued function on a non empty set S , the supremum on the set S

$$\|f\|_S = \sup_{x \in S} |f(x)|$$

If f is a bounded function on S then

$$\sup_{x \in S} |f(x)| = \sup \{ |f(x)| : x \in S \}$$

exists Observe that $|f(x)| \leq \|f\|_S \quad \forall x \in S$

UNIFORM CONVERGENCE IMPLIES POINT WISE CONVERGENCE

From uniform convergence

$$\left| f_n(x) - f(x) \right| \leq \sup_{x \in S} \left| f_n(x) - f(x) \right| \\ = \| f_n - f \|_S$$

So that $f_n \xrightarrow[n \rightarrow \infty]{\text{uniformly on } S} f$

$$\Rightarrow \left| f_n(x) - f(x) \right| \xrightarrow[n \rightarrow \infty]{\text{for each } x \in S} 0$$
$$\Rightarrow f_n \xrightarrow[n \rightarrow \infty]{\text{point wise on } S} f$$

Theorem : Suppose $\{f_n(x)\}_{n \geq 0}$ is a sequence of continuous functions on an interval S . Suppose $f_n(x)$ converges uniformly to $f(x)$ on S . Then the limit function $f(x)$ is also continuous.

We need to establish

$$\forall x, a \in S.$$

$$f(x) \xrightarrow{x \rightarrow a} f(a)$$

PROOF :

Let us start with

$$|f(x) - f(a)|$$

For any $n \geq 0$ i.e., $n = 0, 1, 2, \dots$

$$\begin{aligned} |f(x) - f(a)| &= \left| (f(x) - f_n(x)) + (f_n(x) - f_n(a)) \right. \\ &\quad \left. + (f_n(a) - f(a)) \right| \\ &\leq \left| f(x) - f_n(x) \right| + \left| f_n(x) - f_n(a) \right| \\ &\quad + \left| f_n(a) - f(a) \right| \quad (\because \text{TRIANGLE INEQUALITY}) \end{aligned}$$

$$|f(x) - f(a)| \leq 2 \|f - f_n\|_S + |f_n(a) - f(a)|$$

(

$$|f(x) - f_n(x)| \leq \|f - f_n\|_S$$

$$|f(a) - f_n(a)| \leq \|f - f_n\|_S$$

)

Choose a positive number $\varepsilon > 0$ arbitrary small

$$\|f - f_n\|_S \xrightarrow{n \rightarrow \infty} 0$$

$\therefore \exists N > 0$ for which $\|f_n - f\|_S < \frac{\varepsilon}{3}$

$\forall n \geq N$

Now, $f_N(x)$ is continuous. So, for any choice of $\varepsilon > 0$, there is an interval centered around 'a' so that

$$|f_N(x) - f_N(a)| < \frac{\varepsilon}{3}$$

whenever $x \in$ that interval

Formally, since $f_N(x) \xrightarrow{x \rightarrow a} f_N(a)$
 $\forall \varepsilon > 0$, there is a corresponding $S > 0$ so that whenever $|x-a| < S$

Thus,

$$\left| f(x) - f(a) \right| \leq 2 \left\| f - f_N \right\|_S + \underbrace{\left| f_N(x) - f_N(a) \right|}_{\leq \frac{\epsilon}{3}} = \epsilon$$

$$+ |x - a| < s$$

$$\Rightarrow f(x) \xrightarrow{x \rightarrow a} f(a)$$

$$\Rightarrow f(x) \text{ is } \underline{\text{CONTINUOUS}}$$

Quick Test towards uniform Convergence

If $\{f_n\}_{n \geq 0}$ is a seq. of continuous functions,
the limit function $f(x)$ is 'NOT' continuous
However if $f_n(x)$ converges pointwise to $f(x)$,
 $\Rightarrow f_n(x)$ does not converge uniformly to $f(x)$

Theorem : Suppose $\{f_n(x)\}_{n \geq 0}$ is a seq. of continuous functions which converges uniformly to a cont. function $f(x)$ on a bounded interval $[a, b]$. We

have

$$\lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \int_a^b \lim_{n \rightarrow \infty} f_n(x) dx$$

$$= \int_a^b f(x) dx$$

Proof :

$$\begin{aligned} & \left| \int_a^b f_n(x) dx - \int_a^b f(x) dx \right| \\ &= \left| \int_a^b (f_n(x) - f(x)) dx \right| \\ &\leq \int_a^b |(f_n(x) - f(x))| dx \\ &\leq \int_a^b \|f_n - f\|_S dx = \|f_n - f\|_S \int_a^b 1 \cdot dx \\ &= \|f_n - f\|_S (b-a) \xrightarrow[n \rightarrow \infty]{} 0 \end{aligned}$$

Fourier Series : Properties and notions of convergence

Fourier Series : Over the interval $-\pi \leq x \leq \pi$

$$f(x) = a_0 + \sum_{k=1}^{\infty} a_k \cos(kx) + b_k \sin(kx)$$

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(nx) dx ; \quad b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(nx) dx$$

a_i 's & b_i 's are the Fourier Coeffts.


Consider the function $f(x) = x$ on $-\pi \leq x \leq \pi$

$$f(x) = -f(-x) \quad (\text{odd function})$$

$$\therefore b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} x \sin(kx) dx$$

$$F(x) = \sum_{k=1}^{\infty} \frac{2(-1)^{k+1}}{k} \sin(kx)$$

For this example, $f(x)$ is not 2π periodic ! Let us
form a function \tilde{f} which is a periodic extension of f .




$k = \text{odd multiples}$

$f(x)$ converges to $\tilde{f}(x)$ at points where \tilde{f} is continuous.
 However, we have discontinuity points @ $\pm k\pi$, $k \in \mathbb{Z}; k \neq 0$
 At these points $F(x)$ will converge to $\frac{f(k\pi+) + f(k\pi-)}{2}$

Consider the finite sum

$$S_N(x) = \sum_{k=1}^N \frac{(-1)^{k+1}}{k} \sin(kx)$$



The blips occurring just before and after the points of discontinuity is, called GIBBS EFFECT !

Accuracy of the blips around the discontinuity points gets 'worse'!

The graph of $S_N(x)$ for small finite 'N' say $N=10$ must travel from π to $-\pi$ in a "very short interval".

- 1) The height of the blip is \sim same for large N
- 2) The width gets smaller as N gets larger.

Exercise :

1) Plot $S_N(x)$ for $N = 10, 100, 1000, \dots$
 Observe Gibbs effect.

2) Investigate for a saw tooth wave



$$s(t) = \begin{cases} t & 0 \leq t \leq \frac{\pi}{2} \\ \pi - t & \frac{\pi}{2} \leq t \leq \pi \end{cases}$$

What do you observe?

Theorem : Suppose f is a piecewise continuous function on the interval $a \leq x \leq b$, then

$$\lim_{k \rightarrow \infty} \int_a^b f(x) \cos(kx) dx = \lim_{k \rightarrow \infty} \int_a^b f(x) \sin(kx) dx = 0$$

Proof :



As the frequency $k \uparrow$, $f(x)$ is nearly constant on two adjacent periods of $\sin(kx)$ and $\cos(kx)$. The integral over each such small period is zero over piecewise intervals.

In the case of a differentiable function

$$\int_a^b f(x) \cos(kx) dx = \frac{\sin(kx)}{k} f(x)$$

$$\textcircled{I}: \frac{\sin(kb)f(b) - \sin(ka)f(a)}{k}$$

(Apply L'Hospital's rule)

$$f \quad \int_a^b \frac{\sin(kx)}{k} f'(x) dx$$

↑ \textcircled{II}

\textcircled{III} since $f'(x) = \text{const.}$

Convergence at a point of Continuity

A Fourier series of f converges to f at a point x

if

$$f(x) = a_0 + \lim_{N \rightarrow \infty} \sum_{k=1}^N a_k \cos(kx) + b_k \sin(kx)$$

Theorem : Suppose f is continuous and a 2π periodic function.
Then, at each point ' x ' where $f'(x)$ is defined,
the F.S. of f at x converges to $f(x)$.

Proof:

For a +ve integer N , let

$$S_N(x) = a_0 + \sum_{k=1}^N a_k \cos(kx) + b_k \sin(kx)$$

What we need to show:

$$S_N(x) \xrightarrow[N \rightarrow \infty]{} f(x)$$

for this, let us rewrite S_N in a slightly different way

STEP 1 :

Substituting the Fourier Coeffts,

$$S_N(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) dt + \sum_{k=1}^{N} \left(\int_{-\pi}^{\pi} f(t) \cos kt dt \right) \cos(kx) + \left(\int_{-\pi}^{\pi} f(t) \sin kt dt \right) \sin(kx) \quad \textcircled{1}$$

Using

$$\cos(A - B) = \cos A \cos B + \sin A \sin B \quad \text{in } \textcircled{1},$$

$$S_N(x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) \left(\frac{1}{2} + \sum_{k=1}^{N} \cos[k(t-x)] \right) dt$$

STEP 2 :

Lemma : For any number $u \in [-\pi, \pi]$

$$\frac{1}{2} + \cos u + \cos(2u) + \dots + \cos(Nu) = \begin{cases} \frac{\sin\left(N+\frac{1}{2}\right)u}{2 \sin(u/2)} & u \neq 0 \\ \left(N+\frac{1}{2}\right) & u = 0 \end{cases}$$

Proof :

$$\begin{aligned}
 (e^{jw})^N &= \cos(Nw) + j\sin(Nw) \\
 \frac{1}{2} + \sum_{k=1}^N \cos(kw) &= -\frac{1}{2} + \operatorname{Re} \left\{ \sum_{k=0}^N (e^{jw})^k \right\} \\
 &= -\frac{1}{2} + \operatorname{Re} \left\{ \frac{1 - e^{j(N+1)w}}{1 - e^{jw}} \right\} \\
 &= -\frac{1}{2} + \operatorname{Re} \left\{ \frac{e^{-jw/2} - e^{j(N+\frac{1}{2})w}}{-jw/2} \right\} \\
 &= -\frac{1}{2} + \operatorname{Re} \left\{ \frac{e^{jw/2} - e^{-j(N+\frac{1}{2})w}}{\cos w/2 - j\sin(w/2) - [\cos(N+\frac{1}{2})w + j\sin(N+\frac{1}{2})w]} \right\}
 \end{aligned}$$

$$= -\frac{1}{2} + \frac{\sin\left(N+\frac{1}{2}\right)u + \sin\left(\frac{u}{2}\right)}{2\sin(u/2)}$$

$$= \begin{cases} \frac{\sin\left(N+\frac{1}{2}\right)u}{2\sin(u/2)} & u \neq 0 \\ N + \frac{1}{2} & u = 0 \end{cases}$$

□

STEP 3 :

Let us evaluate the partial sum of the Fourier Series.

$$S_N(x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) \left[\frac{1}{2} + \sum_{k=1}^N \cos[k(t-x)] \right] dt$$
$$= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) \frac{\sin[(N+\frac{1}{2})(t-x)]}{\sin[(t-x)/2]} dt$$

Define

$$P_N(u) = \frac{1}{2\pi} \frac{\sin((N+\frac{1}{2})u)}{\sin(u/2)}$$

$$S_n(x) = \int_{-\pi}^{\pi} f(t) P_n(t-x) dt$$

Let $u = t - x$

$$S_n(x) = \int_{-\pi}^{\pi} f(u+x) \underbrace{P_n(u)}_{\text{"Kernel"}} du$$

(Ponder why the limits on the definite integral remained as is instead of $\pi-x$ & $-\pi-x$)

STEP 4 :

Lemma:

$$\int_{-\pi}^{\pi} P_N(u) du = 1$$

Proof:

Using

$$P_N(u) = \frac{1}{\pi} \left[\frac{1}{2} + \cos u + \cos 2u + \dots + \cos(Nu) \right]$$

$$\Rightarrow \int_{-\pi}^{\pi} P_N(u) du = 1$$

All the $\cos(ku)$ for $k \neq 0$ integrate to zero!

STEP 5 :

Consider $\int_{-\pi}^{\pi} f(u+x) P_N(u) du$ (From Step 3)

From Step 4 Lemma,

$$f(x) = \int_{-\pi}^{\pi} f(x) P_N(u) du$$

We are home if we show

$$\int_{-\pi}^{\pi} [f(u+x) - f(x)] P_N(u) du \xrightarrow[N \rightarrow \infty]{} 0$$

Now,

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{f(u+x) - f(x)}{\sin(u/2)} \sin\left[\left(N + \frac{1}{2}\right)u\right] du$$

Invoking prev. theorem, $\lim_{k \rightarrow \infty} \int_{-\pi}^{\pi} f(x) \sin(kx) \rightarrow 0$
 except @ $u = 0, \pi, -\pi$

$$g(u) = \frac{f(u+x) - f(x)}{\sin(u/2)}$$

is continuous except at $u=0$ over $[-\pi, \pi]$

$$f'(x) = \lim_{u \rightarrow 0} \frac{f(u+x) - f(x)}{u}$$

(f' exists by hypothesis)

Consider

$$\lim_{u \rightarrow 0} \frac{f(u+x) - f(x)}{u} \cdot \frac{\frac{u}{2}}{\sin(\frac{u}{2})} \cdot 2$$
$$= f'(x) \cdot 1 \cdot 2 = 2 f'(x)$$

\equiv
Let us define $g(0) = 2 f'(x)$, $g(u)$ extends across $u=0$ as a cont. function

$$\int_{-\pi}^{\pi} [f(u+x) - f(x)] P_N(u) du \xrightarrow[N \rightarrow \infty]{} 0$$

13

Convergence at a point of discontinuity

Not all functions are continuous and periodic.

If we consider periodic extensions, it may not necessarily result in a continuous function.

Defn:

$$\lim_{h \rightarrow 0^+} f(x-h) = f(x-0) \quad (\text{left limit})$$


$$\lim_{h \rightarrow 0^+} f(x+h) = f(x+0) \quad (\text{right limit})$$

A function f is left differentiable at x if

$$f'(x-0) = \lim_{h \rightarrow 0^-} \frac{f(x+h) - f(x)}{h}$$

Similarly, a function f is right differentiable @ x if

$$f'(x+0) = \lim_{h \rightarrow 0^+} \frac{f(x+h) - f(x)}{h}$$



For $f(x) = x$ over $-\pi \leq x < \pi$ & periodic extensions

$$f(\pi+0) = -\pi \quad f(\pi-0) = \pi$$

$$f'(0) = -1 \quad f'(\pi-0) = 1$$

for $f(x) = \begin{cases} x & 0 \leq x \leq \frac{\pi}{2} \\ \pi-x & \frac{\pi}{2} \leq x \leq \pi \end{cases}$

$f'(x)$ is not differentiable @ $\pi/2$

Theorem : Suppose $f(x)$ is periodic and piecewise continuous-

Suppose x is a point where f is left and right differentiable (but not continuous). The Fourier series of

f at x converges to

$$\frac{f(x+0) + f(x-0)}{2}$$

Proof:

let us slightly deviate from Step 4 of our previous theorem

$$\int_0^{\pi} P_N(u) du = \int_{-\pi}^0 P_N(u) du = \frac{1}{2} \quad \text{--- } ①$$

Recall: $P_N(u) = \frac{1}{2\pi} \frac{\sin\left[\left(N+\frac{1}{2}\right)u\right]}{\sin(u/2)}$ (Even function)
Hence ①

To prove the theorem, we need to show

$$\int_{-\pi}^{\pi} f(u+x) P_N(u) du \xrightarrow[N \rightarrow \infty]{} \frac{f(x+0) + f(x-0)}{2}$$

$$\int_{-\pi}^{\pi} f(u+x) P_N(u) du \xrightarrow[N \rightarrow \infty]{} \frac{f(x+0)}{2} \quad \textcircled{A}$$

$$\int_{-\pi}^0 f(u+x) P_N(u) du \xrightarrow[N \rightarrow \infty]{} \frac{f(x-0)}{2} \quad \textcircled{B}$$

Using the definition of $P_N(u)$

$$\frac{1}{2\pi} \int_0^{\pi} \frac{f(x+u) - f(x+0)}{\sin(u/2)} \sin\left(N + \frac{1}{2}\right) u \ du \xrightarrow[N \rightarrow \infty]{} 0$$

Again

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{f(x+u) - f(x-0)}{\sin(u/2)} \sin\left(N + \frac{1}{2}\right) u \ du \xrightarrow[N \rightarrow \infty]{} 0$$

ASIDE : When we have finite such discontinuities such that the measure on that set $\rightarrow 0$, we are okay with the convergence proof.



Uniform Convergence

Defn : Given a sequence of functions $\{f_n(x)\}$, for a given tolerance $\epsilon > 0 \exists N$ such that $|f_n(x) - f(x)| < \epsilon \forall x$ and $n \geq N$.

Defn : A function is piecewise 'smooth' if it is continuous and its derivative is defined everywhere except possibly at a discrete set of points. Ex: saw tooth waveform



Theorem : The Fourier series of a piecewise smooth
2 π periodic function converges uniformly to $f(x)$
on $[-\pi, \pi]$

Proof : We shall prove this theorem with the assumption
that f is everywhere twice differentiable.

$$\text{Let } f(x) = \sum_n a_n \cos(nx) + b_n \sin(nx)$$

$$f''(x) = \sum_n a_n'' \cos(nx) + b_n'' \sin(nx) \quad \text{(A)}$$

Here $a_n'' = -a_n \cdot n^2$; $b_n'' = -b_n \cdot n^2$

Consider $\sum_{n=1}^{\infty} |a_n| + |b_n| = \sum_{n=1}^{\infty} \frac{|a_n''|}{n^2} + \frac{|b_n''|}{n^2}$ ————— (B)

If f'' is continuous, then a'' & b'' stay bounded by a quantity say ' M ' and ' N '

\therefore (B) Can be written as an inequality

$$\leq \sum_{n=1}^{\infty} \frac{M}{n^2} + \frac{N}{n^2}$$

$$= (M+N) \sum_{n=1}^{\infty} \frac{1}{n^2}$$

To ensure uniform conv., we need another result. $\sum_{n=1}^{\infty} \frac{1}{n^2}$ finite & bounded $\frac{\pi^2}{6}$

Lemma : Suppose $f(x) = a_0 + \sum_{k=1}^{\infty} a_k \cos(kx) + b_k \sin(kx)$

with $\sum_{k=1}^{\infty} |a_k| + |b_k| < \infty$, then

F. S. converges uniformly & absolutely to the function.

Proof : By triangular inequality,

$$|a_k \cos(kx) + b_k \sin(kx)| \leq |a_k| + |b_k|$$

($|\cos(\cdot)| \& |\sin(\cdot)| \leq 1$)

$$\text{Let } S_N(x) = a_0 + \sum_{k=1}^N a_k \cos(kx) + b_k \sin(kx)$$

$$f(x) - S_N(x) = \sum_{k=N+1}^{\infty} a_k \cos(kx) + b_k \sin(kx)$$

$$|f(x) - S_N(x)| \leq \sum_{k=N+1}^{\infty} |a_k| + |b_k|$$

But, with our twice differentiability condition,
 $\sum_{k=N+1}^{\infty} |\overline{a_k}| + |\overline{b_k}| < \infty$

\therefore For a given $\epsilon > 0 \exists N_0 > 0$ so that
 $N > N_0 \Rightarrow \sum_{k=N+1}^{\infty} |a_k| + |b_k| < \epsilon$ irrespective
 of $|x|$

\Rightarrow Uniform Convergence
 Use Lemma, to Conclude the Theorem.

1/1

1/1