

NNSP-1

Homework #5

Instructor: Prof. Shayan Garani Srinivasa

TA: Prayag Gowgi

Solution for 8.5

Given : eigenvalue $\lambda_1 = 1 + \sigma^2$, corresponding eigenvector $q_1 = s$ and $R = ss^T + \sigma^2\mathbf{I}$.
Consider

$$Rq_1 = ss^T s + \sigma^2\mathbf{I}s \quad (1)$$

$$= \mathbf{I}s + \sigma^2\mathbf{I}s \quad (2)$$

$$= (\mathbf{I} + \sigma^2\mathbf{I})s \quad (3)$$

$$= \lambda_1 q_1 \quad (4)$$

Solution for 8.15

Let us center $\bar{\Phi}(x_i)$ using the empirical mean as follows :

$$\bar{\Phi}(x_i) = \Phi(x_i) - \frac{1}{N} \sum_{i=1}^N \Phi(x_i). \quad (5)$$


We define, \bar{K}_{ij} as follows :

$$\bar{K}_{ij} = \langle \bar{\Phi}(x_i), \bar{\Phi}(x_j) \rangle. \quad (6)$$


Substituting equation (5) in equation (6) and simplifying we get

$$\bar{K}_{ij} = K_{ij} - \frac{1}{N} \sum_{m=1}^N \Phi(x_m)^T \Phi(x_j) - \frac{1}{N} \sum_{n=1}^N \Phi(x_i)^T \Phi(x_n) + \frac{1}{N^2} \sum_{m=1}^N \sum_{n=1}^N \Phi(x_m)^T \Phi(x_n) \quad (7)$$

Solution for 8.17



(a) Learning curve of GHA for Lena image.



(b) Learning curve of GHA for pepper image.

FIGURE 1 – Learning curves of GHA

Solution for 8.18

Given :

1. $x_1 \in \mathcal{U}[-1, 1]$.
2. $x_2 = x_1^2 + v$.
3. $v \in \mathcal{N}(0, 0.04)$.
4. Use Kernel Hebbian Algorithm.




FIGURE 2 – Kernel Hebbian algorithm for 2D data.

1. Visually, Figure 2 is similar to the Figure 8.13 in Haykin book.